Pulsar Wind Nebulae

On their Evolution and Diversity

Samar Safi-Harb

Dept of Physics and Astronomy University of Manitoba, Canada samar@physics.umanitoba.ca

IAU Symposium 291 Beijing, 20-24 Aug. 2012



The Most "famous"

that confirmed B&Z's 1934 prediction that neutron stars are born in SNe The brightest and the closest...




Fritz Zwicky

Crab Pulsar NP0532

Staelin & Reifenstein (1968), Lovelace+ (1968), Cocke+1969


Vela PSR0833-45 (89ms) Large+ (1968)

Radio Optical X-rays




Credit: J. Hester (ASU), CXC, HST, NRAO, NSF, NASA

Pulsars are Rotating Neutron Stars with strong B: Magnetized particle wind



(late) Franco Pacini: 1939-2012 (also Gold and Hoyle; Pacini & Salvati, Rees & Gunn)

Pulsar Wind Nebulae






- PWN: Bubbles of relativistic particles inflated by pulsar's relativistic wind

 - √gamma-rays: IC scattering (synchrotron, CMB, IR, starlight), pion decay (hadronic)
- "Calorimeters" for the study of pulsars/Pathfinders for Pulsar Discovery!
- Probes for the Interaction of their relativistic winds with the surrounding: SN ejecta (earlier) or ISM (later): PWN evolution
- Seed the Galaxy with energetic particles and magnetic fields
- Efficient Engines for Cosmic Ray Acceleration up to TeV energies

Chandra: New/Sharp Eyes on PWNe

Jet-Torus sample




Jets/Torus: Numerical Simulations following the time-dependent evolution (e.g. Del Zanna et al.'06, Buccianitni et al.'04; van der Swaluw et al.'04; Blondin, Chevalier, Frieson'01)


Samar Safi-Harb IAUS291, Beijing


PVN variability (movies)

v~fraction of c





PWN Evolution






Chandra Pavlov et al. 2003







"relic" PWNe observed in TeV gamma-rays

van der swaluw 2004 (also Blondin et al. 2001)

Bow-Snock PWN Ram Pressure Confinement: Bow shock+tail (trail) 15 10" 25 <4"→ 26 20 22 10" 3" 6" 29 12 33 👫 30 34 40" 38 30" 36 50" 39 13" 51 25 47 30" 52 50" 15" 49 Kargaltsev & Pavlov 2008 Samar Safi-I

Some Statistics

- As of today, we know of 308 Galactic SNRs (274+34; a new and up-to-date HE SNRcat, which complements Dave Green's catalogue in radio)
- ~90 SNRs contain PWNe or candidates
 - √~35 are HESS sources
 - √~30 lack shells (including the Crab): "naked PWNe"




A New High-Energy (X+g) SNRs catalogue: http://www.physics.umanitoba.ca/snr/SNRcat
Ferrand & Safi-Harb 2012, AdvSpRes
being updated with PWN data
(a dedicated, up-to-date, PWN catalogue)


see poster IAUS291-1-**0206**


The shell is missing in many (Crab-like) PWNe, including the Crab itself!

Chandra pinpoints the blast wave in a "Crab"-like PWN



Evidence for X-ray "Shells" around 2 other PWNe






3C58, XMM-Newton *Gotthelf et al.* 2007


G54.1+0.3, XMM-Newton and Suzaku Bocchino et al. 2010


Not-so-Crab-like

Two unusual Radio Nebulae



- Roughly annular morphologies with inner/outer radii of ~ 1'/10'
- Unlike shell-type SNRs, not limb-brightened ✓ their non-thermal flux falls off away from centre, implying a central engine (pulsar)
- Radio spectrum too steep for a plerion, $\alpha \approx 0.6$, more typical of shells
- Low-frequency (radio) spectral break
- In polarized emission, strong axisymmetry and double-lobed morphology
- See Arzoumanian et al. (2011, 2008).









Radio and X-ray pulsar searches lead to the discovery of a very energetic pulsar! The 2nd most energetic after the Crab pulsar and the fastest known pulsar in our Galaxy!!







P = 24.3 ms

 $P = 4.2 \times 10^{-14} \text{ s s}^{-1}$

 $DM = 427 \text{ pc cm}^{-3}$

 $B \approx 1 \times 10^{12} \text{ G}$

Edot= 1.2×10^{38} erg/s


20h20m45s

20h20m30s

20h20m15

Spectral fits (Chandra)							
Component	Nebula	PSRJ2022					
Model	Power Law	Power Law					
Γ	1.4	1.0 ± 0.2					
n _H (10 ²² cm ⁻²)	$(1.6 \pm 0.3) \times 10^{22} \text{ cm}^{-2}$						
Flux (abs 2–10 keV; ergs cm ⁻² s ⁻¹)	4×10 ⁻¹⁴	5.3×10 ⁻¹³					
Lx (2-10 keV)	5.6x10 ³²	7x10 ³³					


G76.9+1.0




Chandra pinpoints a "pulsar" candidate


- Very hard X-ray spectrum
- Unusually low Lx/Edot (6e-5)
- PWN appears "old"
- PSR appears "young"

Arzoumanian, Gotthelf, Ransom, SSH, Landecker & Kothes 2011, ApJ



G65.7+1.2






Spectral fits (Chandra)


Component	Nebula Pt srce			
Model	Power Law	Blackbody		
Γ / kT_{eff} (MK)	1.6	2.5		
n _H (10 ²² cm ⁻²)	< 0.4			
Flux (unabs 0.5–10 keV; ergs cm ⁻² s ⁻¹)	1.7×10 ⁻¹³	8×10 ^{–14}		
Lx (0.5-10 keV)	3.3x10 ³¹	~10 ³¹		

- •Spectral and energetic modelling suggest 20 kyr age and strong, ~1 mG, PWN field.
- •Implied spin-down luminosity: ~10³⁵ ergs/s
- •No pulsations in Arecibo and XMM-Newton searches.

Another two non-Crab like radio nebulae



- •steep central component $(alpha \sim 0.4 - 0.5)$
- Low-freq. spectral break
- Large, implying evolved?



G63.7+1.1: A PWN interacting with a molecular cloud??

G74.9+1.2: A Relic/Evolved PWN


- $-N_{\rm H} = 1.6 (1.0 2.2) \times 10^{22} \, \rm cm^{-2}$
- X-ray flux = 2.7×10^{-13} erg cm⁻² s⁻¹

- N_H=1.38 (1.21-1.57)x10²² cm⁻²
- ✓ steepens away from point source


Chandra (and XMM) reveal lowluminosity and "offset" X-ray nebulae in collaboration with Roland Kothes

G74.9+1.2

Chandra reveals a compact and an extended "offset" PWN




- Plerion (radio)
- 8'x6' size in the radio (red contours)
- •Low-frequency spectral break, no pulsar
- New sensitive radio studies:
 - √ a larger size of ~16': a linear size of ~28 pc (at 6.1 kpc) and indicating a highly evolved SNR
- ✓ Unusually steep spectral index, $\alpha \approx$ 0.48 (for the central cpt) and a harder component outward



G74.9+1.2


Chandra reveals a compact and an extended "offset" PWN



- Plerion (radio)
- 8'x6' size in the radio (red contours)
- •Low-frequency spectral break, no pulsar
- New sensitive radio studies:
 - √ a larger size of ~16': a linear size of ~28 pc (at 6.1 kpc) and indicating a highly evolved SNR
- ✓ Unusually steep spectral index, $\alpha \approx$ 0.48 (for the central cpt) and a harder component outward

Nebula: cometary morphology


- ✓ neutron star (candidate) moving to the south-east? Offset of ~2' requires a speed of ~175 (km/s)/т₂₀
- ✓ Crushed/relic PWN as e.g. in Vela-X, G327.1-1.1.
- ▶Offset between R, X; steep central radio spectrum
- ▶Bridge connecting them.
- ▶Gamma-ray detection (Aliu et al).



SSH et al. 2011, Matheson, SSH & Kothes 2012

High-B Pulsars (HBPs) in SNRs $\sim 10^{13} < B (Gauss) < \sim 10^{14}$

PSR J1846-0258 in SNR Kes75 P = 0.3 s $B=5 \times 10^{13} G$ Edot=8.3e36 erg/s



PSR J1119-6127 in SNR G292.2-0.5

P=408 ms

B=4.1e13 Gauss~BQED

Edot=2.3e36 erg/s

Age ~1900 yrs

Do HBPs power nebulae?

Samar Safi-Harb I


J1846-0258 in SNR Kes75

(<u>B=5e13 Gauss</u>, Edot=8.3e36 erg/s, Age ~ 725 yrs) powered by both rotation and magnetism




J1846-0258 in SNR Kes75


(B=5e13 Gauss, Edot=8.3e36 erg/s, Age ~ 725 yrs) powered by both rotation and magnetism



J1846-0258 in SNR Kes75

(B=5e13 Gauss, Edot=8.3e36 erg/s, Age ~ 725 yrs) powered by both rotation and magnetism






Ng et al. 2008

high Lx/Edot~0.2-0.3 magnetically powered?

Variability in the PWN following magnetar-like bursting activity (Kumar & SSH 2008; Gavriil et al. 2008; Ng et al. 2008)


J1119-6127 in SNR G292.2-0.5

(<u>B=4.1e13 Gauss</u>, Edot=2.3e36 erg/s, Age ~ 1,900 yrs)






- High pulsed fraction in soft X-ray (Gonzalez+05)
- Compact PWN, Southern Jet
- Power law with a hard photon index
- Energetics can be explained by pulsar's Edot SSH & Kumar 2008



XMM-Newton Energy Image with radio contours (Caswell et al.) overlaid Kumar, SSH & Gonzalez 2012, ApJ

JIII 19-6127 in SNR G292.2-0.5 multi-wavelength view



Kumar, SSH & Gonzalez 2012, ApJ see poster IAUS291-0255

PWNe around other highly magnetized pulsars


THE ASTROPHYSICAL JOURNAL, 703:L41-L45, 2009 September 20
© 2009. The American Astronomical Society. All rights reserved. Printed in the U.S.A.


doi:10.1088/0004-637X/703/1/L41

DISCOVERY OF EXTENDED X-RAY EMISSION AROUND THE HIGHLY MAGNETIC RRAT J1819–1458

N. Rea^{1,2}, M. A. McLaughlin^{3,4,11}, B. M. Gaensler⁵, P. O. Slane⁶, L. Stella⁷, S. P. Reynolds⁸, M. Burgay⁹, G. L. Israel⁷, A. Possenti⁹, and S. Chatterjee¹⁰

see talk later this session by Ascension Camero et al





DRAFT VERSION JULY 30, 2012 Preprint typeset using LATEX style emulateapj v. 5/2/11

see talk in IAUS290 Friday morning

XMM-Newton VIEW OF Swift J1834.9-0846 AND ITS MAGNETAR WIND NEBULA

G. Younes^{1,2}, C. Kouveliotou^{3,2}, O. Kargaltsev⁴, G. G. Pavlov^{5,6}, E. Göğüş⁷, and S. Wachter⁸

Pre-outburst 2005 - Obs. 1


100"

XMMU J183435.3-084443

CXOU J183442.6-084501

Region B

emission, claimed to be a PWN (Bamba and Vink 2009): variable, strongly correlated with AXP's flux: Dust Scattering Halo (Olausen et al. 2011)



X-ray PVNe around high-B NS (6 HBPs + 1 RRAT, ~23 magnetars)

PSR	B _{s,d} (10 ¹³ G)	Edot	PWN extent	Photon index	L _x / Edot	Comment	Reference
JIII9-6127 (see poster 0255, SSH&Kumar)	4.1	2.30E+36	6"x15" 20" jet	. ^{+0.9} -0.7 .4+ ^{0.8} -0.9	5.00E-04	properties similar to RPP's PWNe (Lx/Edot~1e-6-1e-2, gamma~1-2.5)	Gonzalez & SSH 2003 SSH & Kumar 2008
J1846-0258 Crab+magnetar-like (see poster 0228 by Konar)	5	8.30E+36	~40"	~1.9 (1.7-2.2)	~0.2-0.3	variable	Kumar & SSH 2008 Ng et al. 2008
RRAT1819-1458 (see talk by Camero)	5	3.00E+32	13"	3.0+/-1.5	~0.2	see talk (next)	Rea et al. 2009
Swift J1834.9-0846 MWN?	14	2.10E+34	70"-150"	3.5+/-0.6	~0.7	properties similar to RRAT's PWN	Younes et al. 2012
1E 1347.0-3406	22	i.00E+35	45 (2.7)	~3.5	û.û i	most likely dust scattering halo	Bamba & Vink 09 Olausen et al. 2011

Whether PWNe are powered by rotation and/or magnetism, and are associated with other high-B pulsars is still an open question..

Samar Safi-Harb IAUS291, Beijing





Astro-H

(2014)

- **Soft X-ray Spectroscopic** system (SXS, 0.3-12 keV, 3'FOV, ~5eV resolution)
- Soft X-ray Imaging system (0.3-12 keV. 38' FOV)
- Hard X-ray Imager
 (5-80 keV, 9' FOV, 1.7')
 (comparable to NuSTAR)
- Soft Gamma-Ray Detector (50-600 keV)

0.3-600 keV!


searching for thermal emission from synchrotron dominated PWNe (and SNRs) simultaneous modelling of both the thermal & non-thermal plasma in SNR,




Astro-H and PWNe

"Naked" PWNe (Crab-like):

With the SXS, search for the long-sought **missed** thermal emission associated with the blast wave/ shocked ejecta (as hinted e.g. in G21.5-0.9).





Evolved/Crushed PWNe:

With the SXS (+HXD), one can study the interaction between the reverse shock and the (crushed) PWN. Mixed (soft) thermal and (hard) non-thermal X-ray emission detected, e.g. in G327.1-1.1.

Summary

- A long way since the discovery of 1st NS/SNR in 1968!
- Diversity (with still many questions):
 - intrinsic (NS properties, SN progenitor, ambient medium)
 - ✓ growing evidence for shells around Crab-like PWNe
 - √ growing evidence for PWNe around highly magnetized neutron stars
 - Age effect/Evolution (interaction with reverse shock, SNR shell, ambient medium)
 - PWNe as pathfinders for pulsars:
 - ✓ a low Lx does **not** necessarily imply a low Edot.
 - √ look for pulsars in peak of x-ray emission (not necessarily radio)
- Near Future (high-energy):
 - ✓ <u>NuSTAR</u> (June 2012-; hard x-ray emission) <u>Astro-H</u> (2014; thermal plasma in PWNe, broadband spectroscopy). <u>ASTROSAT and HXMT</u> (broadband/hard x-ray spectra)
 - ✓ HESS II (Jul 26-), and other planned gamma-ray missions (CTA)

Thanks to...













Collaborators including Zaven Arzoumanian, Roland Kothes, Tom Landecker..

Contributions by the NS/PWN/SNR community

Thank you IAUS291!